

Optical Coalescence and Applications in Cavity Quantum Electrodynamics

Claudiu Genes,^{1,2,*} André Xuereb,^{3,4} Guido Pupillo,¹ and Aurélien Dantan⁵

¹ISIS (UMR 7006) and IPCMS (UMR 7504), Université de Strasbourg and CNRS, Strasbourg, France

²Institut für Theoretische Physik, Universität Innsbruck, Technikerstrasse 25, A-6020 Innsbruck, Austria

³Centre for Theoretical Atomic, Molecular and Optical Physics,

School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, United Kingdom

⁴Department of Physics, University of Malta, Msida MSD 2080, Malta

⁵QUANTOP, Danish National Research Foundation Center for Quantum Optics,

Department of Physics and Astronomy, University of Aarhus, 8000 Aarhus C, Denmark

(Dated: April 8, 2013)

We investigate a hitherto largely unexplored regime of cavity quantum electrodynamics in which a highly-reflective element positioned between the end-mirrors of a typical Fabry–Pérot resonator strongly modifies the cavity response function, such that two longitudinal modes with different spatial parity are brought close to frequency degeneracy. We examine applications of this generic ‘optical coalescence’ phenomenon for the generation of enhanced photon–phonon nonlinearities in optomechanics and atom–photon nonlinearities in cavity quantum electrodynamics with strongly-coupled emitters.

PACS numbers: 42.50.Pq, 42.50.Ct, 42.50.Wk, 07.60.Ly

A Fabry–Pérot resonator containing a scattering element between its two end-mirrors represents a paradigmatic system for fundamental light–matter interaction studies, as investigated, e.g., in cavity quantum electrodynamics (CQED) [1] and cavity optomechanics [2]. The theoretical description of light in high-finesse cavities, i.e., cavities that possess resonances spaced significantly further apart than their individual widths, is well-established and the modification of the bare cavity response induced by the presence of a scatterer has been exploited in different regimes, distinguished by the scatterer’s reflectivity. In a first regime, a low-reflectivity scatterer, e.g., an atom, is coupled to a single optical mode that spans the entire cavity. This coupling may be used, for instance, to engineer well-controlled and strong interactions between single atoms and single quanta of light for fundamental tests of quantum mechanics, quantum information processing, and metrology [1, 3]. At the opposite end of the scale, the interaction of cavity fields with massive movable scatterers, as studied in optomechanics, allows for a rich variety of tools to measure and control mechanical motion [2]. The case of a harmonically-bound highly-reflective central element, such as a thin, partially transmitting membrane [4–8], is generally described by means of two spatially-separated cavity modes coupled through photon tunneling. Such a system may be used for quantum non-demolition measurements [9] and the observation of jumps in the occupation number of the oscillator [4, 10, 11], tests of the Landau–Zener effect [12], the generation of nonclassical states of motion [13], quantum information processing [14, 15], etc.

In this paper we use the transfer matrix formalism, as used recently to unify these two regimes [16], to go




FIG. 1. (Color online) (a) Optical cavity with a middle reflector of polarizability ζ_m , accommodating odd (red) and even (blue) modes that couple to outside “left” and “right” modes. (b) Cavity transmission spectra over three free spectral ranges, illustrating the strong shift of the odd resonances toward the even ones as the reflectivity is increased. (c) Transmission spectra (solid black curves) for increasing $|\zeta_m|$. The dashed curves denote the “bare” resonances. ‘Mode-pulling’ effects can be seen by comparing the positions of these resonances and the peaks of the transmission. At coalescence (ζ_m^* is defined in the text) one finds a single peak with close to unity transmission over an increased bandwidth. Past this point the maximum transmission is reduced below unity.

beyond the high-reflectivity case as previously investigated and address the situation where the reflectivity of the middle mirror is increased well past that of the cavity mirrors. We unravel new aspects of CQED in the situation where adjacent bare cavity resonances are strongly pulled together, leading eventually to an *optical coalescence* phenomenon, in which pairs of adjacent resonances with different spatial parity become almost degenerate. The interference between these modes modifies the cavity response in a non-trivial fashion. In particular, close to the coalescence point, high transmission can be

* Email address: claudiu.genes@uibk.ac.at

achieved over a bandwidth significantly larger than the bare cavity linewidth. While this mechanism alone could be relevant for practical applications of optical cavities in interferometry and, e.g., the realization of white-light cavities [17], we show that it also has interesting implications for cavity optomechanics as well as CQED with two-level emitters: (i) On the optomechanical side we examine the case of a cavity containing a movable reflector where the cavity field couples quadratically to the mirror motion. Owing to a coalescence-induced pulling of the resonance frequencies, we show that the intrinsic quadratic optomechanical coupling is strongly enhanced as compared to the situations envisaged in standard optomechanical models [2, 4, 5, 16]. (ii) On the CQED side, we show that the dynamics of a weakly-driven two-level quantum emitter strongly coupled to such a cavity differs from that predicted by the standard single-mode Jaynes–Cummings (SMJC) model. In the good-cavity limit, when the emitter’s dissipation rate is larger than the cavity field decay rate, the system can operate as a photon turnstile device, with strong antibunching and bunching observed in the cavity transmission and reflection spectrum, respectively, over a large bandwidth around the emitter’s resonance frequency. This provides an alternative topology to microtoroid photon turnstile devices, which involve scattering between quasi-degenerate counter-propagating optical modes [18]. Overall, our results show that optical coalescence could provide an alternative route towards achieving strong optomechanical nonlinearities [11, 12, 15, 19, 20] or the dynamical control of light at the quantum level [18, 21].

Model.—Let us consider a cavity of length L consisting of two mirrors with polarizability $\zeta < 0$, related to the amplitude reflectivity through $r = -i\zeta/(1 - i\zeta)$, and a middle reflector with polarizability ζ_m , cf. Fig. 1(a). We assume that the mirrors are lossless (i.e., described by real polarizabilities) and that the middle reflector has a thickness smaller than the wavelengths considered, themselves much smaller than L . We make use of the transfer matrix formalism for one-dimensional scatterers to solve the Helmholtz equation [16, 22], and compute the transmission function of the system for general values of ζ and ζ_m . For an empty cavity with sufficiently good mirrors the transmission spectrum consists of well-separated Lorentzian peaks with frequencies given by $\omega_n = (c/L)[(n-1)\pi + \cos^{-1}(\zeta/\sqrt{1+\zeta^2})]$ ($n = 1, 2, \dots$) and linewidth $\kappa = c/(2L|\zeta|\sqrt{1+\zeta^2})$. In the presence of the middle reflector, adjacent resonance frequencies are shifted closer together by an amount that depends on its position and polarizability [Fig. 1(b)]. When the reflector lies exactly at the center of the cavity, as shown in Fig. 1(a), the peaks quantified by an even mode number (‘e’, frequency ω_e) are shifted strongly towards the odd-numbered (‘o’, ω_o) peaks:

$$\omega_o - \omega_e \approx \frac{c}{L} \tan^{-1} [2\zeta_m/(\zeta_m^2 - 1)] \equiv 2\delta.$$

For modest middle-mirror reflectivities the transmission peaks have non-overlapping Lorentzian profiles, which

ensures that they indeed occur at the cavity resonances [see Fig. 1(c)]. However, as $|\zeta_m|$ is increased, the frequency separation between ω_o and ω_e can become comparable to their linewidth and the two modes begin to interfere with each other. In this regime the peaks of the transmission are no longer located at the cavity resonances but at $\tilde{\omega}_e = \frac{c}{L}[(2n)\pi - \epsilon_-]$ and $\tilde{\omega}_o = \frac{c}{L}[(2n+1)\pi - \epsilon_+]$ where

$$\cos \epsilon_{\pm} = \frac{\zeta_m(2\zeta^2+1)(\zeta\zeta_m-1) \pm (\zeta+\zeta_m)\sqrt{4\zeta^2(\zeta^2+1)-\zeta_m^2}}{2\zeta(\zeta^2+1)(\zeta_m^2+1)}.$$

The interesting behavior happens around the point where $\tilde{\omega}_e = \tilde{\omega}_o$, which we dub *optical coalescence* and which is characterized by the merging together of the two Lorentzian profiles [see the illustration of the mechanism in Fig. 1(c)] thereby giving rise to a qualitatively different cavity response. The threshold to this behavior occurs at $\delta = \kappa$ or, in terms of the polarizabilities, when

$$\zeta_m = \zeta_m^* \equiv 2\zeta\sqrt{\zeta^2 + 1}.$$

As $|\zeta_m|$ is further increased past coalescence, the transmission attains only one peak, whose value rapidly decreases. A striking feature of the coalescence regime is the increased transmission bandwidth. This is reminiscent of white-light cavities familiar in the field of interferometric gravitational-wave detection [17], which allow the detector to benefit from significant intracavity laser power over large bandwidths.

At this point, we note that the criterion $\zeta_m \sim \zeta_m^*$ for observing coalescence-related effects is rather demanding on the quality of the middle reflector, which has to be markedly more reflective than the end-mirrors. However, this condition may be significantly alleviated through the use of a multi-layered reflector [23–25]. Indeed, the effective collective polarizability of a multi-layered array scales approximately exponentially with the number of layers [22, 25]. This yields an effective coalescence threshold $|\zeta_m^*| \approx \sqrt[N]{\zeta^2/2^{N-2}}$ and implies that each element of the middle reflector only needs to be as good as the end-mirrors for $N = 2$ – or even markedly less for $N > 2$ – in order to achieve the coalescence condition.

Simple two-mode model.—Let us now construct a simple two-mode Hamiltonian that provides a basis for the quantum-mechanical applications of coalescence in the context of CQED. As illustrated in Fig. 1(a) adjacent resonances have opposite parities, i.e., if the field profile of one resonance is odd, that of the next is even. Our cavity is patently double-sided, and therefore each mode couples to two infinite baths of harmonic oscillators, one on each side of the cavity $b_{l,r}(\omega)$. Standard techniques and the Markovian assumption [26] allow us to define input (output) fields $b_{l,r}^{\text{in(out)}}(t)$ as appropriate Fourier transforms of $b_{l,r}(\omega)$. A simple argument based on coordinate inversion shows that the even (odd) mode of the pair couples to an even (odd) combination of bath modes. We define two cavity modes $a_{o,e}$ with corresponding frequencies




FIG. 2. (Color online) Mode-pulling behavior of adjacent cavity modes ($\zeta = -10$, $\zeta_m = -196.6$). Shown are two adjacent resonances as a function of middle-mirror displacement about the center of the cavity. For perfect cavity mirrors, i.e., zero linewidth, one obtains the resonances shown by the dotted curves. Imperfect cavity mirrors cause an interference effect that pulls adjacent modes closer together (solid curves), resulting in a sharper avoided crossing, and enhanced quadratic optomechanical coupling.

$\omega_{o,e} = \omega \pm \delta$, such that the free Hamiltonian is simply

$$H_0 = (\omega - \delta)a_o^\dagger a_o + (\omega + \delta)a_e^\dagger a_e.$$

We include the input noise terms for $a_{o,e}$ as the two linear combinations $[b_1^{\text{in}}(t) \pm b_2^{\text{in}}(t)]/\sqrt{2}$. Assuming driving through a given mirror at a frequency Ω , we can calculate the cavity transmission at the other mirror under steady-state conditions, obtaining

$$T(\Omega) = \left| \frac{\kappa}{\kappa + i(\omega - \delta - \Omega)} - \frac{\kappa}{\kappa + i(\omega + \delta - \Omega)} \right|^2.$$

This simple expression, in excellent agreement with the spectra of Fig. 1(c), obtained from the full numerics, clearly shows the interference between the two Lorentzians as one approaches coalescence.

Nonlinear optomechanics.—Let us now consider a first application of optical coalescence to the field of optomechanics, by including a mechanical degree of freedom (displacement x) for the middle mirror. When the equilibrium position of the mirror is at the centre of the cavity, the photon-phonon coupling is quadratic in x , corresponding to a dependence of the cavity resonance frequency on x^2 . Such a quadratic optomechanical coupling has been proposed, e.g., to perform quantum non-demolition measurements (QND) of the phonon number operator [9], possibly resolving single-phonon quantum jumps [5, 10, 11]. For a typical cavity operating far from coalescence, the resonance frequency can be written $\omega(x) \approx \omega + G_2^{(0)}x^2$ (where ω is the resonance at rest) and the quadratic coupling strength $G_2^{(0)} = -2\omega^2/(cL)\zeta_m$, which scales linearly with the reflector polarizability [5, 16]. Membrane-optomechanics experiments [4, 5, 8] usually operate in the regime $|\zeta_m| \ll |\zeta|$, and, although progress in making highly reflective membranes is ongoing [7, 27], this quadratic coupling remains typically much weaker than the linear coupling. In the following we exploit the coalescence mechanism

to demonstrate a scaling of the quadratic coupling with ζ_m beyond this linear case. To this end, we start with our two-mode model described by H_0 to which we add a displacement-induced tunneling term, governed by the tunneling strength $g_m x$:

$$H = H_0 + g_m x (a_o^\dagger a_e + a_o a_e^\dagger). \quad (1)$$

This Hamiltonian has been previously treated (for example in Refs. [11, 20, 23]) and reproduces the transfer matrix results in a quasi-static picture. The transmission peaks occur at $\omega \pm \Omega_\pm(x)$, with $\Omega_\pm = \pm\sqrt{\delta^2 + (g_m x)^2 - \kappa^2}$, and agree with the exact numerically calculated $\tilde{\omega}_{o(e)}$. Comparison with Ref. [16] provides an expression for the tunneling parameter appearing in Eq. (1):

$$g_m = (2\omega/L)\{(\zeta_m/2)\tan^{-1}[2\zeta_m/(\zeta_m^2 - 1)]\}^{1/2}.$$

In the limit of small displacements, $g_m|x| \ll \sqrt{\delta^2 - \kappa^2}$, one obtains the quadratic optomechanical coupling strength

$$G_2(\zeta_m) = \frac{2\zeta^2}{\sqrt{\zeta_m^2 - \zeta_m^2}} G_2^{(0)}.$$

In the two-fold limit of good cavity mirrors ($|\zeta| \gg 1$) and away from coalescence ($|\zeta_m| \ll \zeta^2$), this expression reduces to the coupling $G_2(\zeta_m) \rightarrow G_2^{(0)}$. As one approaches coalescence, however, one observes a dramatic increase in the coupling strength. The physical mechanism at the basis of this enhancement is the coalescence-induced mode-pulling mechanism illustrated in Fig. 2, where the presence of a nearby mode ‘pulls’ a second mode towards the former. An analogy can be drawn between the mechanism we explore here, which amplifies the quadratic coupling through an interference between two longitudinal modes, and the use of multiple *transverse* modes [5] to achieve the same. However, as can be seen from the expression for $\Omega_\pm$, the displacement region over which this enhancement is obtained becomes smaller, and ultimately vanishes, as one approaches coalescence. One can easily show that the quadratic expansion is valid as long as $G_2(\zeta_m)/G_2^{(0)} \ll 2/(\eta|\zeta_m|)$, where $\eta = 2\pi|x|/\lambda$ is the Lamb-Dicke parameter. For typical membrane-based optomechanical experiments experiencing zero-point fluctuations over $\sim \text{fm}$ length-scales and for $\sim 1 \mu\text{m}$ wavelengths, the enhancement can still be of several orders of magnitude. For a stack of two $\sim 100 \text{ ng}$ membranes with frequency $2\pi \times 100 \text{ kHz}$ and $\zeta_m \approx -10$, the enhancement reaches $\sim 10^6$ for ground state-cooled membranes and $\sim 10^3$ for membranes at 4 K. We stress that the enhancement found in this work purely arises from a full treatment of the *static* properties of the system and further note that the application of the Hamiltonian in Eq. (1) to typical optomechanical situations with mechanical resonators vibrating at some frequency ω_m can give rise to rich dynamical effects. In the good-cavity limit $\omega_m \gg \kappa$, one expects the fast mechanical




FIG. 3. (Color online) (a) Single-excitation subspace of the dressed state energy levels for a strongly-coupled two-level atom interacting with weakly pumped cavity modes. See text for the definitions of the kets. (b) Left: Spectra for the anti-symmetric (upper) and symmetric (lower) modes around the atomic resonance. Right: Corresponding values of $g^{(2)}(0)$ as function of the detuning. Values below the dotted lines denote antibunching. Δ is defined as the difference between the driving frequency and ω . ($g = 10\sqrt{2}\kappa$, $\gamma = 5\kappa$, and $\delta = \kappa$, photon pumping rate 0.03κ .)

oscillator to mediate strong interactions between the two optical modes, which can be used, e.g., for achieving enhanced nonlinearities [11, 15] or displacement sensitivity [20]. Such ‘dynamical’ effects are expected to give rise to interesting behavior around the coalescence regime and will be examined elsewhere.

Cavity QED.—We now turn to the application of optical coalescence to cavity QED with emitters. We focus on the strong-coupling regime of CQED in which the intrinsic nonlinearity of a saturable two-level emitter is manifest [18, 21]. Let us assume for simplicity that a two-level emitter is placed into the half-cavity where the two modes have identical spatial profiles, i.e., on the left in the case of the system illustrated in Fig. 1(a). The emitter therefore couples to these modes with an equal coupling rate $g/\sqrt{2}$. The emitter resonance frequency ω is assumed to lie mid-way between the two cavity mode frequencies at $\omega \pm \delta$. The coherent dynamics of the system are described by the free Hamiltonians H_0 for the fields and $\omega\sigma_+\sigma_-$ for the emitter, supplemented by the Jaynes–Cummings interaction Hamiltonian:

$$H_{\text{JC}} = \frac{g}{\sqrt{2}}(a_o^\dagger\sigma_- + a_o\sigma_+) + \frac{g}{\sqrt{2}}(a_e^\dagger\sigma_- + a_e\sigma_+).$$

The non-unitary dynamics are described via the associated master equation, where for a general decay process at rate Γ with operator O the associated Liouvillian is $\mathcal{L}(\rho) = (\Gamma/2)(2O\rho O^\dagger - OO^\dagger\rho - \rho O^\dagger O)$. The incoherent baths for our particular systems are described by rates $\{\gamma, \kappa, \kappa\}$, respectively, corresponding to the collapse operators $\{\sigma_-, a_e, a_o\}$. We define as well the symmetric and antisymmetric field operators $a_s = (a_o + a_e)/\sqrt{2}$ and $a_a = (a_o - a_e)/\sqrt{2}$.

These dynamics can be solved numerically for different driving conditions of the cavity, but in order to draw a parallel with the strong-coupling regime of the SMJC model, we diagonalize the Hamiltonian in the single-excitation (weak driving) limit and look at the dissipation in the dressed-state basis. The energy levels, as shown in Fig. 3(a), are $\epsilon_\pm = \omega \pm \sqrt{g^2 + \delta^2}$ and $\epsilon_0 = \omega$. The associated eigenstates are obtained by applying to the ground state of the system $|0\rangle = |g00\rangle$ the Hermitian conjugates of the following three operators:

$$p_\pm = \frac{1}{\sqrt{2}}(\sigma_- \pm a_s + \frac{\delta}{g}a_a) \text{ and } p_0 = a_a - \frac{\delta}{g}\sigma_- ,$$

to obtain the states $|\psi_\pm\rangle = p_\pm^\dagger|0\rangle$ and $|\psi_0\rangle = p_0^\dagger|0\rangle$. In the good-cavity limit $\gamma \gg \kappa, \delta$, the states of the system are simply two *atom-polaritons* showing the typical vacuum Rabi splitting, supplemented with a *light-polariton* that contains an entangled two-mode field state $(|g10\rangle - |g01\rangle)/\sqrt{2}$. The dissipation in the polariton basis in the same limit shows the expected SMJC harmonic averaging of the decay rates for the atom-polaritons: the decay of the two atom-polaritons is governed by $\mathcal{L}_\pm(\rho)$ at a rate $(\kappa + \gamma)/2$, while the dissipative evolution of the light-polariton described by the Liouvillian $\mathcal{L}_0(\rho)$ is decoupled from the other polaritons and occurs at the rate κ , as illustrated in Fig. 3(a).

The existence of a light-polariton arises from the coalescence-induced mode interference and clearly differs from the SMJC. We compute numerically the zero-time second-order correlation function, $g^{(2)}(0)$, of the output symmetric and antisymmetric modes for a strongly-coupled emitter $(g, \gamma) = (10\sqrt{2}\kappa, 5\kappa)$ at coalescence ($\delta = \kappa$), as shown in Fig. 3(b). While the symmetric mode spectrum resembles a typical SMJC spectrum and displays bunching [$g^{(2)}(0) > 0$] at atomic resonance, the antisymmetric mode exhibits a three-peak spectrum as well as strong antibunching around atomic resonance. For a cavity pumped on one side, antibunching and bunching would then be observed in transmission and reflection, respectively; the system would act as a photon turnstile device in which the emitter regulates the transmission of photons one-by-one. In contrast to photon blockade scenarios relying on the anharmonicity of the Jaynes–Cummings energy levels [21] or nonlinear cavities [28], the dynamical control of the photon transmission here comes from the atom-mediated interference between two cavity modes. An interesting parallel can be made with the experiments of Ref. [18], making use of an atom coupled to a microtoroidal resonator in which backward- and forward-propagating modes are coupled on account of scattering losses. A key difference is the possibility of the system we discussed to operate in the good-cavity limit, which results in a blockade bandwidth of ca. $\gamma \gg \kappa$, rather than κ , as illustrated in Fig. 3(b). We note that coalescence-induced interference between dressed states at stronger driving could give rise to further interesting atom–photon dynamics.

Concluding remarks.—We have explored a fundamental effect in the quantum electrodynamical description of a

cavity with a partially-transmitting mirror inside it, in the limit where this mirror is significantly better than the cavity mirrors themselves, and have shown its potential for achieving significantly stronger photon–phonon coupling in the field of optomechanics, and strong atom–photon nonlinearities in CQED.

We acknowledge support from the Austrian Science Fund

(FWF): P24968-N27 and an STSM grant from the COST Action MP1006 “Fundamental problems in quantum physics” (C.G.), the Royal Commission for the Exhibition of 1851 (A.X.), the Université de Strasbourg through Labex NIE, IDEX, and EOARD (G.P.), and the Danish Council for Independent Research under the Sapere Aude program (A.D.).

-
- [1] P. R. Berman, *Cavity Quantum Electrodynamics*, Adv. At. Mol. Opt. Phys. Series (Academic Press, 1994); S. Haroche and J.-M. Raimond, *Exploring The Quantum: Atoms, Cavities, And Photons* (Oxford University Press, 2006); H. J. Kimble, *Nature* **453**, 1023 (2008).
 - [2] M. Aspelmeyer, T. J. Kippenberg, and F. Marquardt, e-print (2013), [arXiv:1303.0733](#).
 - [3] G. Khitrova, H. M. Gibbs, M. Kira, S. W. Koch, and A. Scherer, *Nat. Phys.* **2**, 81 (2006); A. J. Shields, *Nat. Photonics* **1**, 215 (2007); S. M. Girvin, M. H. Devoret, and R. J. Schoelkopf, *Phys. Scr.* **2009**, 014012 (2009).
 - [4] J. D. Thompson, B. M. Zwickl, A. M. Jayich, F. Marquardt, S. M. Girvin, and J. G. E. Harris, *Nature* **452**, 72 (2008).
 - [5] J. C. Sankey, C. Yang, B. M. Zwickl, A. M. Jayich, and J. G. E. Harris, *Nat. Phys.* **6**, 707 (2010).
 - [6] D. J. Wilson, C. A. Regal, S. B. Papp, and H. J. Kimble, *Phys. Rev. Lett.* **103**, 207204 (2009); K. Usami, A. Naesby, T. Bagci, B. Melholt Nielsen, J. Liu, S. Stobbe, P. Lodahl, and E. S. Polzik, *Nat. Phys.* **8**, 168 (2012).
 - [7] U. Kemiktarak, M. Metcalfe, M. Durand, and J. Lawall, *Appl. Phys. Lett.* **100**, 061124 (2012); U. Kemiktarak, M. Durand, M. Metcalfe, and J. Lawall, *New J. Phys.* **14**, 125010 (2012).
 - [8] M. Karuza, M. Galassi, C. Biancofiore, C. Molinelli, R. Natali, P. Tombesi, G. D. Giuseppe, and D. Vitali, *J. Opt.* **15**, 025704 (2013); T. P. Purdy, R. W. Peterson, and C. A. Regal, *Science* **339**, 801 (2013).
 - [9] V. B. Braginsky, Y. I. Vorontsov, and K. S. Thorne, *Science* **209**, 547 (1980).
 - [10] A. A. Clerk, M. H. Devoret, S. M. Girvin, F. Marquardt, and R. J. Schoelkopf, *Rev. Mod. Phys.* **82**, 1155 (2010).
 - [11] M. Ludwig, A. H. Safavi-Naeini, O. Painter, and F. Marquardt, *Phys. Rev. Lett.* **109**, 063601 (2012).
 - [12] G. Heinrich, J. G. E. Harris, and F. Marquardt, *Phys. Rev. A* **81**, 011801 (2010); N. Miladinovic, F. Hasan, N. Chisholm, I. E. Linnington, E. A. Hinds, and D. H. J. O’Dell, *Phys. Rev. A* **84**, 043822 (2011).
 - [13] S. Bose, K. Jacobs, and P. L. Knight, *Phys. Rev. A* **56**, 4175 (1997); D. Kleckner, I. Pikovski, E. Jeffrey, L. Ament, E. Eliel, J. van den Brink, and D. Bouwmeester, *New J. Phys.* **10**, 095020 (2008); M. Paternostro, *Phys. Rev. Lett.* **106**, 183601 (2011); L. F. Buchmann, L. Zhang, A. Chiruvelli, and P. Meystre, *Phys. Rev. Lett.* **108**, 210403 (2012).
 - [14] K. Stannigel, P. Rabl, A. S. Sørensen, P. Zoller, and M. D. Lukin, *Phys. Rev. Lett.* **105**, 220501 (2010).
 - [15] K. Stannigel, P. Komar, S. J. M. Habraken, S. D. Bennett, M. D. Lukin, P. Zoller, and P. Rabl, *Phys. Rev. Lett.* **109**, 013603 (2012).
 - [16] A. Xuereb and P. Domokos, *New J. Phys.* **14**, 095027 (2012).
 - [17] A. Wicht, K. Danzmann, M. Fleischhauer, M. Scully, G. Müller, and R. H. Rinkleff, *Opt. Commun.* **134**, 431 (1997); G. S. Pati, M. Salit, K. Salit, and M. S. Shahriar, *Phys. Rev. Lett.* **99**, 133601 (2007); A. A. Savchenkov, A. B. Matsko, and L. Maleki, *Opt. Lett.* **31**, 92 (2006).
 - [18] B. Dayan, A. S. Parkins, T. Aoki, E. P. Ostby, K. J. Vahala, and H. J. Kimble, *Science* **319**, 1062 (2008).
 - [19] H. Miao, S. Danilishin, T. Corbitt, and Y. Chen, *Phys. Rev. Lett.* **103**, 100402 (2009).
 - [20] S. Basiri-Esfahani, U. Akram, and G. J. Milburn, *New J. Phys.* **14**, 085017 (2012); H. Seok, L. F. Buchmann, S. Singh, and P. Meystre, *Phys. Rev. A* **86**, 063829 (2012); P. Kómár, S. D. Bennett, K. Stannigel, S. J. M. Habraken, P. Rabl, P. Zoller, and M. D. Lukin, *Phys. Rev. A* **87**, 013839 (2013); X. Xu and J. M. Taylor, e-print (2013), [arXiv:1303.7469](#).
 - [21] K. M. Birnbaum, A. Boca, R. Miller, A. D. Boozer, T. E. Northup, and H. J. Kimble, *Nature* **436**, 87 (2005); I. Schuster, A. Kubanek, A. Fuhrmanek, T. Puppe, P. W. H. Pinkse, K. Murr, and G. Rempe, *Nat. Phys.* **4**, 382 (2008); A. Kubanek, A. Ourjoumtsev, I. Schuster, M. Koch, P. W. H. Pinkse, K. Murr, and G. Rempe, *Phys. Rev. Lett.* **101**, 203602 (2008); J. M. Fink, M. Goppl, M. Baur, R. Bianchetti, P. J. Leek, A. Blais, and A. Wallraff, *Nature* **454**, 315 (2008); L. S. Bishop, J. M. Chow, J. Koch, A. A. Houck, M. H. Devoret, E. Thuneberg, S. M. Girvin, and R. J. Schoelkopf, *Nat. Phys.* **5**, 105 (2009).
 - [22] I. H. Deutsch, R. J. C. Spreeuw, S. L. Rolston, and W. D. Phillips, *Phys. Rev. A* **52**, 1394 (1995).
 - [23] M. Bhattacharya, H. Uys, and P. Meystre, *Phys. Rev. A* **77**, 033819 (2008).
 - [24] M. J. Hartmann and M. B. Plenio, *Phys. Rev. Lett.* **101**, 200503 (2008).
 - [25] A. Xuereb, C. Genes, and A. Dantan, *Phys. Rev. Lett.* **109**, 223601 (2012).
 - [26] C. W. Gardiner and P. Zoller, *Quantum Noise*, 3rd ed. (Springer, 2004).
 - [27] C. H. Bui, J. Zheng, S. W. Hoch, L. Y. T. Lee, J. G. E. Harris, and C. W. Wong, *Appl. Phys. Lett.* **100**, 021110 (2012).
 - [28] A. Imamoglu, H. Schmidt, G. Woods, and M. Deutsch, *Phys. Rev. Lett.* **79**, 1467 (1997); P. Grangier, D. F. Walls, and K. M. Gheri, *Phys. Rev. Lett.* **81**, 2833 (1998).